

# Earth and Space Science



## RESEARCH ARTICLE

10.1029/2021EA002059

†Deceased.

Yun Lin and Yoshihide Takano these authors contributed equally to this work.

### Key Points:

- Hydroclimatic variables in the Upper Colorado River Basin show stronger annual trends in March than other months in early spring
- Thinner clouds spur extra warming and thereby additional runoff decline in March
- Intensified large-scale subsidence leads to dryness and cloud suppression in March

### Supporting Information:

Supporting Information may be found in the online version of this article.

### Correspondence to:

Y. Lin,  
yunlin@ucla.edu

### Citation:

Lin, Y., Takano, Y., Gu, Y., Wang, J., Zhao, B., Liou, K.-N., & Fu, R. (2022). Investigation of springtime cloud influence on regional climate and its implication in runoff decline in Upper Colorado River Basin. *Earth and Space Science*, 9, e2021EA002059. <https://doi.org/10.1029/2021EA002059>

Received 30 SEP 2021

Accepted 12 DEC 2021

### Author Contributions:

**Conceptualization:** Yun Lin, Yoshihide Takano, Yu Gu

**Formal analysis:** Yun Lin, Yoshihide Takano

**Investigation:** Yun Lin, Yoshihide Takano







**Methodology:** Yun Lin, Yoshihide Takano, Yu Gu

**Project Administration:** Rong Fu  
**Supervision:** Yu Gu, Kuo-Nan Liou, Rong Fu

© 2021 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

## Investigation of Springtime Cloud Influence on Regional Climate and Its Implication in Runoff Decline in Upper Colorado River Basin

Yun Lin<sup>1</sup> , Yoshihide Takano<sup>1</sup> , Yu Gu<sup>1</sup> , Jingyu Wang<sup>2</sup> , Bin Zhao<sup>1,3</sup> , Kuo-Nan Liou<sup>1,†</sup>, and Rong Fu<sup>1</sup> 

<sup>1</sup>Department of Atmospheric and Oceanic Sciences, Joint Institute for Regional Earth System Science and Engineering, University of California, Los Angeles, CA, USA, <sup>2</sup>National Institute of Education, Nanyang Technological University, Singapore, Singapore, <sup>3</sup>School of Environment and State Key Joint Laboratory of Environment Simulation and Pollution Control, Tsinghua University, Beijing, China

**Abstract** The subseasonal features of the annual trends of runoff and other associated hydroclimatic variables in the upper Colorado River basin (UCRB) are examined using multiple data sets from in situ observations, reanalysis, and modeling for early spring (February, March, and April), given that about 58% of annual mean runoff decline from 1980 to 2018 stem from its decreases in the three months. Our analysis suggests that the strong annual trends of hydroclimatic variables in March are more statistically significant than other two months. While recent observational studies attribute the decline of runoff for either annual total or cool and warm seasons to regional warming and precipitation decrease, we suggested, for the first time, that a larger decreasing trend of the runoff in March is caused by the declining cloud optical depth which induces further decrease in precipitation and additional increase in temperature on top of climatic warming. The extra warming can reduce available water resource in the basin likely by enhancing evaporation in March. The recent cloud suppression likely results from stronger subsidence and larger moisture flux divergence over southwestern United States because of abnormal circulation patterns in varying climate, in turn leading to drier atmosphere which is unfavorable for cloud formation/development over the UCRB region. The cloud influence on the runoff in March in the UCRB revealed in this study implies the importance of understanding subseasonal variations of hydroclimate in the changing climate, as well as a need of future studies on the response of circulation patterns to climate change at subseasonal level and its implication on local hydroclimate.

## 1. Introduction

Runoff of the upper Colorado River basin (UCRB) is vital to water supply in the United States Southwest and California, but it is especially vulnerable to climate change because of the snowmelt-dominated water supply in the basin (Hock et al., 2019). The recent decline of the UCRB runoff has been widely related to precipitation deficits and rising temperatures under anthropogenic climate change, with the relative importance of different drivers varying with time periods examined and methodologies used (Hoerling et al., 2019; McCabe et al., 2017; Udall & Overpeck, 2017; Vano et al., 2012; Vano & Lettenmaier, 2014; Woodhouse et al., 2016; Xiao et al., 2018). Previous estimations based on instrumental records have documented that most of annual variations of streamflow are explained by precipitation variability in river basins across the western U.S., suggesting the dominant role of precipitation anomalies relative to temperature changes in determining runoff interannual variability (Gleick, 1986; Karl & Riebsame, 1989; Nash & Gleick, 1991; Wigley & Jones, 1985). For example, it is documented that precipitation deficit is the major cause of the midcentury drought in 1950s (Udall & Overpeck, 2017). On the other hand, temperature warming tends to either moderate or exacerbate the influence of precipitation anomalies, overall playing a secondary role in runoff interannual variability in the past (Woodhouse et al., 2016). The contributions to the annual runoff reduction due to the temperature rising can be about 5–10%, according to studies of Vano et al. (2012) over 1975–2005 and Udall and Overpeck (2017) over 2000–2014. However, at a longer term (e.g., a 100-year period between 1916 and 2014), Xiao et al. (2018) found that more than half (53%) of the runoff decreasing trend can be attributed to the warming through hydrologic modeling estimation. They also noted that the precipitation slightly increased (i.e., 1.4%) over this period but with the streamflow decline of 16.5%. In addition, the warming interannual trend over the UCRB is getting pronounced since 1980s (Dawadi & Ahmad, 2013; McCabe et al., 2017) and more frequent warmer years with lower flow since 1988 has been reported (Woodhouse

**Validation:** Yun Lin, Jingyu Wang, Bin Zhao

**Visualization:** Yun Lin, Yoshihide Takano

**Writing – original draft:** Yun Lin, Yoshihide Takano

**Writing – review & editing:** Yun Lin, Yoshihide Takano, Yu Gu, Jingyu Wang, Bin Zhao, Kuo-Nan Liou, Rong Fu


et al., 2016), both of which suggests an increasingly importance of rising temperature in more recent years in controlling the response of runoff. For example, both observational and modeling studies have confirmed that the very important role of rising temperature during the ongoing post-Millennium Drought relative to its counterpart in the midcentury drought (Udall & Overpeck, 2017; Xiao et al., 2018). Soil moisture is another influential factor for runoff, but its impact on runoff is at lesser degree than that of temperature and precipitation (Hamlet et al., 2007; Seneviratne et al., 2010; Woodhouse et al., 2016).

Given the possible increasing influence of climate change on runoff, modeling efforts have been made in the projections of future changes in temperature and precipitation, as well as their impacts on runoff response. Although the sensitivity of streamflow to precipitation (percent runoff change per percent precipitation change) in future projections is less disputable (Hobbins & Barsugli, 2020), the runoff reduction attributable to regional warming (i.e., temperature sensitivity of runoff to temperature, defined by percent discharge change per degree Celsius of warming) shows large variations, ranging from <10% to 45% by 2050 (Vano & Lettenmaier, 2014). The uncertainty is particularly originated from the great discrepancy between empirical-based regressions and hydrologic modeling simulations used to estimate streamflow sensitivity (McCabe et al., 2017; Milly et al., 2018; Nowak et al., 2012; Vano et al., 2012; Vano & Lettenmaier, 2014). Obviously, the current efforts still cannot reconcile each other in streamflow sensitivity estimations. To counter this, more comprehensive process understanding, rigorous synthesis of observations, and modeling works with various model frameworks and full range of temperature perturbation inputs are required. One such study carried out recently by Milly and Dunne (2020) has estimated individual contributions of various processes related to climatic warming to the UCRB discharge change, suggesting that the total annual decrease is about 9.3% °C<sup>-1</sup> and the primary driver of runoff loss is the snow-albedo feedback under the warming climate.

Alternatively, other studies argue that the recent declines of the UCRB runoff could at least partially be attributed to the effect of dust on snowpack (Painter et al., 2010), teleconnections with tropical Pacific sea surface temperature (Lehner et al., 2018; Zhao et al., 2021), the direct effects of rising CO<sub>2</sub> concentration on the balance of long-wave radiation and on water-use efficiency of vegetation, as well as land-cover modifications due to wildfires and insect outbreaks (Hobbins & Barsugli, 2020). However, all these possible drivers still need further more accurate estimations, particularly together with the direct effect of global warming.

On the other hand, the response of streamflow to temperature change in Western United States is characterized with strong subannual variations (Ban et al., 2020; Das et al., 2011; Vano et al., 2015; Vano & Lettenmaier, 2014), and may represent another important source of uncertainties when estimating the sensitivity of annual runoff to regional warming. The examination of subannual variations particularly focuses on warm (April to September) and cool (October to March) seasons. For example, Ban et al. (2020) find that there are asymmetrical responses of annual and seasonal streamflow changes to the warming of warm and cool seasons in river basins like the UCRB in Western United States. In addition to the distinct responses of warm and cool seasons, there exist subseasonal variations of annual changing patterns for hydroclimatic variables in the UCRB. For instance (Miller & Piechota, 2008), report that the annual trend signs of temperature, precipitation, and streamflow in the Colorado River Basin highly vary by month. However, the subseasonal features of hydroclimate in response to regional warming are still rarely touched, and, as a consequence, their driving factors are not fully recognized yet.

In this study, we will combine multiple data sets from in situ observations, reanalysis, and modeling for the UCRB and perform statistical analysis to understand the hydroclimate in the basin at monthly scale, especially in early spring when snowmelt and runoff mostly occur. We aim to comprehensively assess if there are parameters over global warming-induced changes in temperature and precipitation (as well as the related teleconnections) may affect the runoff and other associated hydroclimatic condition in the UCRB. We select the period of 1980–2018 for study because the warming over the UCRB is pronounced since 1980s (Dawadi & Ahmad, 2013; McCabe et al., 2017) and the signature of temperature rising in the runoff interannual trend appears more significant over this period relative to prior, evident in the fact that there is enhanced frequency of warmer year with lower flow (Woodhouse et al., 2016). In addition, our focus on the most recent decades may help to better understand the recent relationships among hydroclimatic variables under global warming, which will assist in accurate modeling projections of future runoff response in the UCRB.



**Figure 1.** Map of upper Colorado River basin (UCRB). A red frame indicates the investigated domain: 35°–43°N, 105.7°–111°W.

## 2. Material and Methodology

For the data collection and analysis, we focus on the UCRB with a domain of 35°–43°N, 105.7°–111°W, which is indicated with a red frame in Figure 1. The monthly hydroclimatic variables and their associated variables used in this study include surface runoff, temperature, precipitation, soil moisture, snow water equivalent (SWE), evapotranspiration, surface net radiation flux, cloud optical depth (COD), and surface and planetary albedo (ALB) for the period 1980–2018 (except 1999–2018 for SWE based on its availability). Among these variables examined, temperature, precipitation, and soil moisture are the drivers of runoff response in the changing climate suggested by most previous studies. SWE, surface net radiation flux, and albedo parameters are closely related to water supply in snow-fed basins, and can be directly modulated by hydroclimatic conditions like temperature and precipitation (e.g., Hamlet et al., 2007; McCabe et al., 2017; Milly & Dunne, 2020). The COD is selected to probe the possible effects due to cloudiness changes on shortwave radiation attenuation, and thus surface thermodynamic conditions (Hamlet et al., 2007; Vano et al., 2012; Udall & Overpeck, 2017). Surface runoff data were extracted from Global Land Data Assimilation System (GLDAS) simulations with Common Land Model (CLM), which has been proven to be able to realistically capture the seasonal and interannual variability (Dai et al., 2003; Zaitchik et al., 2010). In order to further validate that

the data adopted in this study is suitable for the annual trend analysis for the UCRB over the study period, the GLDAS-CLM simulation of runoff was validated against USGS streamflow gauge measurements at Lees Ferry station which measures the amount of water that flows out of UCRB and simulations from North American Land Data Assimilation System (NLDAS) with a finer horizontal spacing 0.125° by 0.125° than GLDAS (i.e., 1° by 1°). NLDAS ensemble mean was derived based on available simulations using three land surface models (i.e., Noah, Variable Infiltration Capacity (VIC), and Mosaic) and used for comparison. The GLDAS-CLM simulation is certainly smaller than USGS gauge measurements because it simulates only surface runoff only while the gauge measurements include both surface runoff and base flow. It is found that the GLDAS-CLM simulation better catches the observed runoff interannual variations and its decline trend than does NLDAS ensemble mean (Figure S1a), and shows a good correlation with USGS measurements (Figure S1b). Therefore, we chose GLDAS simulated runoff for its annual trend analysis. The station surface temperature was obtained through World Meteorological Organization KNMI Climate Explorer, the data of which is provided by Global Historical Climatology Network and the full list of stations in the network can be found in Table S2. Precipitation and soil moisture were from Climate Prediction Center (CPC), following the recommendation by Smith and Kummerow (2013); the reanalysis data of surface shortwave (SW) and longwave (LW) radiative fluxes, COD, and ALB were extracted from Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA2). Given that MERRA2 has assimilated numerous meteorological observations like temperature and wind profiles (Gelaro et al., 2017), the variables extracted from MERRA2 are likely reliable. As the two key variables in this study (i.e., COD and temperature) extracted from MERRA2, we particularly evaluated them against available observations using the Moderate Resolution Imaging Spectroradiometer (MODIS) level 3 product (i.e., MOD08M3) for COD and Parameter elevation Regression on Independent Slopes Model (PRISM) data for temperature. It is found that MERRA2 COD in general captures the observed interannual variations in each month (Figure S2a). However, the MODIS observation is only available from 2003 to 2018, but the decline trend of COD can be better recognized at a longer term (e.g., 1980–2018 in this study). Therefore, the use of MERRA2 data in this study appears to be reasonable. In addition, since MERRA2 well reproduces the interannual variation and subseasonal temporal feature as shown in PRISM, it is proper for us to conduct annual trend analysis at monthly level based on MERRA2 temperature (Figure S2b). SWE was from Canadian Meteorological Center (CMC). To verify the CMC SWE data, a comparison was made between CMC and 4-km University of Arizona snow data product, which suggests that the two data sets show very similar interannual variations during the overlapping period (i.e., 1998–2018), but the SWE values from University of Arizona are much larger than those from CMC (Figure S3). It is found that the CMC SWE values are comparable to previous studies, for example, Hamlet et al. (2007). Therefore, we adopted CMC data in this study for analysis. Evapotranspiration data were derived as the ensemble means

of three sets of available GLDAS simulations using different land surface models, including Catchment Land Surface Model (CLSM), Noah, and VIC. Other variables used for large-scale circulation pattern analysis include three-dimensional 500 hPa geopotential height, temperature, relative humidity, water vapor, and wind speeds are all extracted from MERRA2. For the gridded data set, each variable is averaged over the grid points within the UCRB. By doing so, the basinwide mean COD can represent the cloudiness in the UCRB to some extent because the average is done over all the grid elements unconditionally in the basin (including cloudy and noncloudy elements). The nongridded temperature data is averaged over all stations inside the domain. All the data sets used in this study are also listed in Table S1 with the information on spatial resolutions and accessible online sources.

Yearly time series for monthly mean of each hydroclimatic variable were subjected to statistical analysis using Mann-Kendall test in an attempt to detect the subseasonal characteristics of their annual trends between 1980 and 2018. The Mann-Kendall test is a nonparametric test and does not require the normality of the data set, and thereby is popular for the trend analysis in climatologic and in hydrologic time series (Mavromatis & Stathis, 2011; Yue & Wang, 2004). Another advantage of the Mann-Kendall test is its insensitivity to abrupt breaks caused by inhomogeneous time series. Throughout the analysis, the Mann-Kendall trend significance test was accessed at the significance level  $\alpha$  at 0.05 and/or 0.1. The Theil-Sen slopes were derived for denoting the trending intensity.


Pearson's correlation analysis was performed to evaluate the associations between two interested hydroclimatic variables. Since partial correlation coefficient is a measure of the dependence between two variables when the influence of possible controlling variables is removed (Engström & Ekman, 2010; Hardle & Simar, 2015; Johnson & Wichern, 2007; Zhao et al., 2018), we also conducted partial correlation analysis following (Engström & Ekman, 2010) for runoff with temperature and precipitation as the influence of other possible drivers were removed. More details about partial correlation calculation can be found in Text S1 in Supporting Information S1. In addition, the relative contribution of individual predictor (e.g., temperature, precipitation, and so on) to variations of runoff was estimated by performing multiple linear regression analysis. The detailed description about multiple linear regression is in Texts S2–S4 in Supporting Information S1.

### 3. Results

The overall trends are illustrated in the time series of water-year (defined as the 12-month period from October 1 for any given year through September 30 of the following year) annual runoff, temperature, and precipitation in the UCRB (Figure S4). The relative variation of the annual runoff is comparable to the natural flow of Colorado River at Lees Ferry reported by Woodhouse et al. (2016). The UCRB runoff shows a strong decreasing trend from 1980 to 2018, whereas temperature exhibits an increasing trend and no significant trend is on precipitation, suggesting that the decline of runoff is primarily caused by rising temperature. Our results appear more consistent with Xiao et al.'s (2018) study on the period of 1916–2014, in which more than half of the runoff reduction is attributed to temperature warming, but different from others (Gleick, 1986; Karl & Riebsame, 1989; McCabe et al., 2017; Nash & Gleick, 1991; Udall & Overpeck, 2017; Vano et al., 2012; Wigley & Jones, 1985), in which the negative precipitation anomalies are the main driver of runoff decline, with a minor role of temperature changes. The two reasons below could be responsible for the inconsistency between our results with others: First, our study focus on the most recent period with more pronounced warming than that in other studies (e.g., 1975–2005 in Vano et al. (2012)), and thereby the temperature is likely more influential in our study. This is also revealed by Woodhouse et al. (2016). Second, the great interannual variation of precipitation over the studying period likely leads to a difficulty to recognize the signature of precipitation changing in the runoff interannual trend. However, the changes in precipitation did contribute to the runoff variation during this period, which is evident in the fact that the low runoff since 2000 (i.e., the post-Millennium Drought) was well associated with the low level of precipitation over the same period (Figure S4).

To understand the seasonal variations, we examine multiyear monthly means of runoff, temperature, precipitation, and SWE averaged over the UCRB (Figure 2a). The primary peak of runoff occurs in March, while the maximum of SWE occurs in February, 1 month earlier than the primary peak of runoff. On average, the runoff in February, March, and April together accounts for ~51% of the annual total runoff. Figure 2b shows annual anomalies with respect to the 1980–2018 mean. It is found that the annual total anomalies from 1980 to 2018 are mainly attributed to the changes in early spring, that is, February, March, and April. Based on the annual trend (i.e., the linear fitting in Figure S4), the difference in annual total runoff between 1980 and 2018 is estimated






**Figure 2.** (a) Multiyear monthly mean of runoff, temperature, precipitation, and snow water equivalent (SWE) averaged over the UCRB. The primary runoff peak occurs in March. (b) The time series of annual UCRB runoff curve (left axis) and its anomalies in stack bars (right axis) with respective to (1980–2018) mean. The contributions of February, March, and April to annual total runoff anomaly are marked as green, red, and blue bars, respectively. Only the months with runoff change with same sign as the annual total anomaly, that is, positive monthly contributions, are shown in stack columns.

about  $-34.7 \text{ kg m}^{-2}$ , of which February, March, and April contribute about  $-7.0$ ,  $-7.6$ , and  $-5.4 \text{ kg m}^{-2}$ , respectively. Therefore, there is totally  $\sim 58\%$  annual mean loss of runoff from the changes in the three months of early spring. As such, our remaining analysis will focus on the runoff interannual trends and the driving factors in these three months.

Figure 3a shows that the decreasing trends on runoff in all the 3 months pass the Mann-Kendall significance test ( $\alpha = 0.05$ ), with annual mean rates of  $-0.16$ ,  $-0.20$ , and  $-0.12 \text{ kg m}^{-2} \text{ yr}^{-1}$  for February, March, and April, respectively. As suggested by previous studies, the strong trends on runoff can be attributed to the variations in temperature, precipitation and soil moisture, which is also evident in Figures 3b–3d in this study. The temperature rising (Figure 3b), most likely due to global warming, together with the decreasing of soil moisture (Figure 3d) are responsible for the significant annual loss of runoff for all the 3 months. As suggested by previous studies, for example, Huntington and Billmire (2014); Milly and Dunne (2020); Pascolini-Campbell et al. (2021), the increased temperature can enhance evaporation and reduce soil moisture and lead to reduction of runoff. Figures 3d and 3h show that the evaporation/soil moisture in all the 3 months show an increasing/decreasing trend in the past decades. However, the runoff decline is moderated by precipitation increase in February and April, leading to weaker runoff trends in these 2 months than March.

It is noteworthy that the runoff in March declines more dramatically than February and April, as indicated by a faster decreasing rate in March than the other 2 months (Figure 3a). Meanwhile, Figure 3b shows that the warming rate of temperature in March (i.e.,  $0.05 \text{ }^\circ\text{C yr}^{-1}$ ) is several-time higher than those in February (i.e.,  $0.02 \text{ }^\circ\text{C yr}^{-1}$ ) and April (i.e.,  $0.01 \text{ }^\circ\text{C yr}^{-1}$ ). Moreover, the March precipitation shows a decreasing trend with a



**Figure 3.** Time series of monthly (a) runoff, (b) surface temperature, (c) precipitation, (d) soil moisture, (e) surface net radiative flux, (f) snow water equivalent (SWE), (g) cloud optical depth (COD), and (h) evapotranspiration rate in the UCRB in February (green dash), March (red dash), and April (blue dash), and their annual trend fittings (thick solid lines) during 1980–2018. Numbers in each panel denote the Theil-Sen slopes of linear fittings, with the asterisks denoting those trends pass the Mann-Kendall trend significance test at the  $\alpha$  of 0.05 (\*) or 0.10 (\*\*) level.

relatively large rate of  $0.4 \text{ mm yr}^{-1}$  (Figure 3c), opposite to the slight increasing trends in February and April. Clearly, the combined effect of faster temperature warming and precipitation reduction is responsible for the stronger runoff decreasing in March. Although runoff is highly correlated with soil moisture in all the 3 months (Table S3), we can rule out the possibility of soil moisture variability as the primary driving factor for the most significant annual loss in runoff in March, since both the magnitude and the decreasing trend in soil moisture are highly consistent among the 3 months as shown in Figure 3d.

Another distinct feature of March from February and April is that all the annual trends of the runoff-related hydroclimatic variables examined here, including surface temperature, precipitation, soil moisture, surface net radiative flux, SWE, and evapotranspiration rate, pass the Mann-Kendall trend significance test, which is not the case (except evapotranspiration rate and soil moisture) for February and April. This further confirms that the hydroclimatic condition in March in the UCRB is distinct from February and April, likely stemming from a very different underlying mechanism leading to the runoff decline in March with respect to the other 2 months.

Given that clouds can regulate the precipitation distribution and provide variable radiative effects, they are an essential link in the global and regional water cycle as well as the energy budget. Therefore, it is very likely that cloud variations may also play a role in subseasonal hydroclimatic response. We find that the COD in March presents a very strong, statistically significant decreasing trend, while no such trends are found in February and April (Figure 3g). The decreasing trend in March is also manifested by the strong positive correlation between

COD and runoff with a relatively high correlation coefficient of 0.40 (Table S3). Since the COD in March is also closely correlated with temperature ( $R = -0.40$ ) and precipitation ( $R = 0.87$ ), the changes in clouds may contribute to the faster decline of runoff in March by modulating temperature and precipitation. In the following discussions, we will further scrutinize this possible influence on the runoff from cloud.

The relative importance of cloud effect on runoff is evaluated by conducting partial correlation analysis to obtain partial correlation coefficients between runoff and temperature (i.e.,  $R_{(RO,Temp)\bullet COD}$ ) and precipitation (i.e.,  $R_{(RO,Preci)\bullet COD}$ ) when the influence of COD is excluded. The results together with regular correlation coefficients ( $R_{RO,Temp}$  and  $R_{RO,Preci}$ , respectively) are reported in Table 1. When COD is controlled in regressions for February, the positive

**Table 1**  
Correlation Coefficient and Partial Correlation Coefficient Between UCRB Variables of Runoff (RO), Temperature, Precipitation, and Cloud Optical Depth (COD) (1980–2018)

	$R_{RO,Temp}$	$R_{(RO,Temp)\bullet COD}$	$R_{RO,Preci}$	$R_{(RO,Preci)\bullet COD}$
February	-0.28	-0.36	0.44	0.26
March	-0.31	-0.19	0.38	0.06
April	-0.54	-0.52	0.19	0.08

Note.  $R_{A,B}$  is correlation coefficient between A and B.  $R_{(A,B)\bullet C}$  is partial correlation coefficient between A and B while controlling the effect of C.


correlation between runoff and precipitation ( $R_{RO,Preci} = 0.44$ ) becomes weaker ( $R_{(RO,Preci) \bullet COD} = 0.26$ ), but the negative correlation between runoff and temperature ( $R_{RO,Temp} = -0.28$ ) gets stronger ( $R_{(RO,Temp) \bullet COD} = -0.36$ ). As such, the opposite effects of cloud on temperature and precipitation cancel each other out, and overall the cloud does not contribute much to the decline of runoff. In April, both  $R_{RO,Temp}$  ( $-0.54$ ) and  $R_{RO,Preci}$  ( $0.19$ ) becomes smaller ( $-0.52$  and  $0.08$ , respectively) due to the removal of COD influence, but the changes are relatively small, indicating that the changes in COD might be not the primary driver of the runoff decreasing trend in April. In March, the absolute values of  $R_{RO,Temp}$  ( $-0.31$ ) and  $R_{RO,Preci}$  ( $0.38$ ) are both substantially reduced to  $-0.19$  and  $0.06$ , respectively, when COD is controlled, supporting that that the cloud influence is one of the main drivers of the decreasing trend of runoff through causing additional increase in temperature and decrease in precipitation. Further multiple linear regression analysis, as described in Text S3 in Supporting Information S1, reveals that COD changes contribute by 25% of the total explainable variation in runoff by regression in March. In contrast, only 6% and 4% of the explainable variations in runoff can be attributed to COD changes in February and April, respectively (Texts S2 and S4 in Supporting Information S1). The multiple regression analysis obtained here also supports more predominant contribution of cloud influence to runoff in March than in February and April, crossvalidating the conclusion based on the partial correlation analysis in Table 1.

The correlation analysis shows that the COD has a strong negative correlation with the surface net (SW + LW) radiation flux, with correlation coefficients of  $-0.45$ ,  $-0.81$ , and  $-0.73$  for February, March, and April, respectively (Table S3). In other words, the thinner cloud leads to less attenuation of solar radiation and therefore more incoming SW radiation reaching surface. In past several decades, the COD in the UCRB shows a significant decline trend in March, and correspondingly, the surface net radiation flux presents an obvious increasing trend in the same month (Figure 3e). More surface radiation results in additional warming which is evidenced by its strong positive correlation with surface temperature ( $R = 0.63$ ). As a result, the temperature in March warms faster than other 2 months (Figure 3b), indicating an additional warming due to COD reduction on top of climate warming in March. The enhancement in surface temperature induced by COD reduction can further melt more snow, consistent with the strong decreasing trend on SWE in March (Figure 3f). A warmer temperature can reduce runoff through other pathways, for example, by spurring evapotranspiration and sublimation and decreasing soil moisture, which has been extensively demonstrated (Huntington & Billmire, 2014; Milly & Dunne, 2020; Pascolini-Campbell et al., 2021). Figure 3h suggests that the annual trend in the evaporation in March is stronger with a larger increasing slope ( $0.14$ ) than other 2 months (i.e.,  $0.10$  and  $0.12$ , respectively), which is consistent with its greater warming. The relationship analysis shows a moderate correlation between evapotranspiration rate and temperature or surface net radiative flux in March ( $R = 0.36$  or  $0.37$ , respectively) but there is no significant correlation found in February and April, also suggesting that the likely important role of the extra warming in March in reducing available water resource in the basin by enhancing evapotranspiration.

On the other hand, strong associations between COD and precipitation are found for all the 3 months ( $R > 0.78$ ), consistent with other studies, for example, (Halimi et al., 2017; Mishra, 2019). The close co-variation of the two parameters is understandable because COD is proportionated to cloud thickness as a first approximation and the precipitation probability increases as the cloud thickness increases (Sakakibara, 1978). Therefore, the magnitude and the underlying mechanism of the subseasonal variation of precipitation in the UCRB can be inferred via the understanding of how the monthly COD changes in past several decades in the same region.


Then another question arises why the March COD in the UCRB is significantly reduced in the past decades while no such trends are found in February and April? This is likely related to the changes in thermodynamic environment in which clouds form. Given that relative humidity (RH) is closely related to cloud formation (Lamquin et al., 2009; Walcek, 1994), here we pay appreciable attention to the RH annual trend and its controlling factors to understand the influence of thermodynamic condition on clouds. A strong relationship between COD and RH indeed exists in our study ( $R = 0.8$ , Figure 4a). In addition, we find that the column-mean RH (1,000–300 hPa) during 1980–2018 presents a decreasing trend in all the 3 months, but only the trend in March is statistically significant, with a much larger slope (i.e.,  $0.25\% \text{ yr}^{-1}$ ) than the other 2 months (i.e.,  $0.04\% \text{ yr}^{-1}$  and  $0.10\% \text{ yr}^{-1}$  for February and April, respectively, see Figure 4b). Meanwhile, the column-mean air temperature warms at a much higher rate (i.e.,  $0.07 \text{ K yr}^{-1}$ ) in March than the other 2 months (i.e.,  $0.01 \text{ K yr}^{-1}$  and  $0.03 \text{ K yr}^{-1}$  for February and April, respectively, see Figure 4c). In addition to the warming and drying in March occurring in the entire atmosphere column, we separately examined the trends of RH and temperature at different levels as shown in Figure S5, where the strong decreasing/increasing trends of RH/temperature are found at all atmospheric levels.






**Figure 4.** (a) Linear regression between cloud optical depth (COD) and relative humidity (RH) with 95% confidence limits (gray shading) and 95% prediction limits (dash lines), and time series of column-mean (1,000–300 hPa) (b) RH and (c) temperature in UCRB in February (green), March (red), and April (blue), and their trends (thick solid lines) for the period 1980–2018. Numbers in (b, c) denote the Theil-Sen slopes of linear fittings, with the asterisks denoting that the linear trend passes the Mann-Kendall trend significance test at the  $\alpha$  of 0.05 level. Green, red and blue colors in all panels represent February, March, and April, respectively.

To understand the physics of the distinct subseasonal variations of RH and temperature in the UCRB, we further examine monthly changes of large-scale flows in past several decades and their implications for local hydroclimatic condition. Figure 5 depicts the mean differences of midatmospheric flow, vertical velocity, and moisture transportation between the last 10 years (i.e., 2009–2018) and the first 10 years (i.e., 1980–1989) of the study period. The positive 500-hPa geopotential height difference is found in March over the western US region including the UCRB, suggesting that this region is more likely controlled by ridges aloft in recent years. The circulation shows distinct patterns among different months (Figure 5a), as the strong positive anomaly is located west of the UCRB in February, then it propagates right above the UCRB in March, and finally further propagates to the east coast in April. This pattern is also evident in the annual trend analysis, in which the mean 500-hPa geopotential height over the UCRB increases at a much greater rate in March than the other 2 months (Figure 6a). The ridge control




**Figure 5.** The difference between the last 10 years (2009–2018) and the first 10 years (1980–1989) in (a) geopotential height at 500 hPa ( $Z_{500}$ , shading) superimposed with wind vectors at same level, (b) vertical velocity ( $\omega$ , shading) along latitude-height cross section over UCRB superimposed with vectors of (V component of wind,  $-\omega$ ), and (c) vertically integrated moisture flux divergence (1,000–300 hPa, shading) over UCRB domain superimposed with vertically integrated moisture flux (1,000–300 hPa, vectors). The green boxes in (a) mark the UCRB domain, which is domain used for analysis in (b, c). From left to right in each panel are February, March and April. The UCRB domain-mean moisture flux divergence for each month is marked in (c), which are yellow-highlighted. All the data are obtained from MERRA2.





**Figure 6.** Time series of domain-mean (a) geopotential height at 500 mb and (b) vertically integrated moisture flux divergence over UCRB in February (green dash), March (red dash), and April (blue dash), and their trend (thick solid lines) for the period 1980–2018.

is manifested by strong subsidence as shown in Figure 5b middle panel, in contrast to the slight subsidence/ascent patterns found in February/April. The dominant subsidence control in March further constrains the vertical transport of moisture from lower levels, supporting the increased temperature and reduced RH in Figures 4b and 4c. Regarding the moisture supply advected from regions outside of the UCRB, we find that the vertically integrated (1,000–300 hPa) moisture flux divergence in the UCRB is enhanced in all the 3 months, indicating that the UCRB is the moisture source rather than the sink in its neighboring area. Moreover, the domain-averaged enhancement in March (i.e.,  $0.8 \times 10^{-5} \text{ kg m}^{-2} \text{ s}^{-1}$ ) is greater than the other 2 months (i.e.,  $0.3 \times 10^{-5} \text{ kg m}^{-2} \text{ s}^{-1}$  and  $0.5 \times 10^{-5} \text{ kg m}^{-2} \text{ s}^{-1}$  for February and April, respectively, see Figure 5c). This is in line with the annual trends, showing moisture flux divergence in March increases with a larger slope than that in February and April (Figure 6b). The stronger moisture flux divergence in March reduces the external moist supply and further contributes to the atmosphere drying over the UCRB in addition to the subsidence contribution.



**Figure 7.** Flow diagram for the causative relationships among large-scale circulation characteristics, cloud optical depth, and hydroclimatic variables (including runoff) in the UCRB in early spring of 1980–2018 (except 1999–2018 for SWE) under varying climate. Red and blue arrows denote increasing and decreasing trends, respectively. The key factors cloud optical depth, temperature, precipitation, and runoff are highlighted in bold. The background gray giant arrow suggests that all the changes are originated from abnormal circulation patterns under changing climate. For February and April, the hydroclimatic response is encompassed in the dashed box, as it starts from variations of surface temperature and precipitation which are likely resulted from climatic warming alone.

#### 4. Discussion

The flow diagram in Figure 7 summarizes the causative linkage between large-scale air flow perturbations and cloudiness change in March in the UCRB, as well as how this change leads to hydroclimate responses, particularly to the runoff decline. Figure 7 shows that, in March of past several decades, the stronger subsidence and larger moisture divergence were revealed in the UCRB likely due to stronger or more persistent the middle-level ridges controlling southwestern US, which in turn results in dryness in the middle and lower atmosphere over the basin. As a result, the cloud formation/development is suppressed, as indicated by the significant reduction in COD, which accompanies less precipitation and solar radiation attenuation. The increased surface solar radiation contributes additional warming at the surface, further reducing the available water supply in the basin for runoff by enhancing evaporation and sublimation, causing drought (e.g., soil moisture deficit), or redistributing SWE to earlier months.

Our explanation by linking the more significant runoff decline in March to the cloud influence can be an important complement to the current understanding of the hydroclimate response in varying climate, in which the runoff decreasing in basins like the UCRB has been wildly attributed to temperature increasing and precipitation decreasing under global warming (Milly & Dunne, 2020; Udall & Overpeck, 2017; Woodhouse et al., 2016). To probe the possible subseasonal responses of hydroclimate to global warming, we

performed additional analysis using the High Resolution WRF Simulations of the Current and Future Climate of North America data set (Liu et al., 2017), in which two sets of 13-year ensemble simulations with and without of pseudoglobal warming (PGW) were conducted. In comparison to the control simulations, the simulated surface temperature with PGW is enhanced by 4–5 °C in all the 3 months, and more precipitation is produced in both February and March (Figures S6a and S6b). Moreover, the total condensate water path in March increases by 10% with PGW relative to the control case (Figure S6c), which is opposite to the decreasing trend of COD suggested above. This suggests that, at least from modeling perspective, the direct effect of climatic warming alone is not able to interpret the distinct subseasonal variations of the annual trends on hydroclimatic variables in early spring.

A recent study by Milly and Dunne (2020) reveals that the decline of Colorado River flow stemming from enhanced evapotranspiration under global warming is mainly driven by snow-albedo feedback. Our analysis also suggests a weak decreasing trend of surface albedo (Figure S7) along with the continuous snow losses annually (i.e., SWE reduction in Figure 3f) in the 3 months, supporting that the climatic warming-induced snow-albedo feedback may play a minor role in the water reduction in the UCRB. It is worth noting that the planetary albedo in March exhibits a statistically significant decreasing trend with a much larger slope than the other 2 months (Figure S8), which cannot be explained by the interannual variations of surface albedo since the similar decreasing trends on surface albedo are found in all the 3 months. In fact, the great decrease in planetary albedo in March in past decades coincides with the remarkable reduction in COD (Figure 3g). In other words, the greater annual decline of the runoff in March is closely associated with the larger suppression of cloud formation/development. As such, the cloud influence could be regarded as an additional factor driving the decline of runoff in March on top of global warming, that is, the COD reduction leads to extra warming at surface, which in turn induce excess evapotranspiration and thereby excess losses of water resource in the basin in March. In contrast, the decreasing of the runoff in February and April appears mainly due to the global warming effect alone.

Although the larger runoff decline in the UCRB in March is not directly associated with the general global warming effect, the stronger subsidence in March leading to the drier and warmer atmosphere which is unfavorable for cloud formation is likely associated with the abnormal circulation patterns in past decades under climate changes. For example, Kirk et al. (2017) found that strong ridge patterns along with similar dry patterns have been increasingly common in the spring and early summer, contributing to the early 21st century drought across western North America including the UCRB. Several hypotheses have been proposed to elaborate the dynamical mechanisms of the enhanced likelihood of ridge frequency and persistency over the western United States (Burgdorf et al., 2019; Kirk & Schmidlin, 2018; Namias, 1983). One of them argues that the Arctic amplification because of rapid Arctic warming contributes to slower zonal propagation of weather patterns, which is attributable to weakened zonal winds, allowing ridges lingering over certain regions for longer time (Francis & Vavrus, 2015; Kirk & Schmidlin, 2018). In addition, the reduced zonal winds in the midlatitude may result in less contributions from orographic lift to precipitation production in the western United States. Another possible mechanism is related to the poleward shift of subtropical jet stream that is associated with tropical expansion under global warming (Burgdorf et al., 2019). The poleward-shifted jet accompanies the establishment of ridge control over the western United States, which contributes to the subsidence condition over the region and thereby suppressing cloud formation. However, all these hypotheses mentioned here still need further in-depth studies from both observational and modeling perspectives in the future.

## 5. Conclusions

Using multiple data sets from in situ observations, reanalysis, and modeling, surface and atmospheric variables in the UCRB are analyzed in terms of trend, correlation coefficient, and multiyear monthly mean to examine the recent hydroclimate changes, particularly for the declining runoff in early spring (i.e., February, March and April), given that more than half of the annual mean runoff decline from 1980 to 2018 results from decreases in early spring and that the maximum runoff normally occurs in March. Our analysis shows that the statistically significant annual trends of hydroclimatic parameters are much stronger in March than February and April. This is because, besides the temperature increase associated with global warming, the decrease of COD contributes to the larger decreasing trend of the UCRB runoff in March through its effects on both temperature and precipitation. Multiple linear regression analysis indicates that 25% of the variation of runoff can be explained by COD reduction in March, suggesting that the changes in cloud can exert an additional effect on the runoff on the top of global warming. The recent cloud reduction is likely attributed to the changes in large-scale atmospheric

circulations in a changing climate. The abnormal circulation patterns in March tend to cause stronger or persistent ridge control over southwestern United States, contributing to strong subsidence and moisture divergence over the UCRB, thereby suppressing cloud formation. The cloud influence proposed in this study is able to fill the knowledge gap in understanding subseasonal variations of hydroclimate in the UCRB. Our analysis also indicates that further studies may be required to understand the abnormal circulation patterns in the changing climate at subseasonal level.

## Data Availability Statement

The Global Land Data Assimilation System (GLDAS) runoff data were used in this work, which can be downloaded from [https://disc.gsfc.nasa.gov/datasets/GLDAS\\_CLM10\\_M\\_V001/summary](https://disc.gsfc.nasa.gov/datasets/GLDAS_CLM10_M_V001/summary). Evapotranspiration data from GLDAS are obtained from <https://disc.gsfc.nasa.gov>. KNMI temperature data can be accessed via <https://climexp.knmi.nl/selectstation.cgi?id=someone@somewhere>. The precipitation and soil moisture data were downloaded from Climate Prediction Center (CPC) via <https://www.esrl.noaa.gov/psd/data/gridded/data.unified.html>. Snow water equivalent (SWE) was obtained from Canadian Meteorological Centre (CMC), which can be accessed via <https://nsidc.org/data/nsidc-0447>. All other parameters including SW + LW, COD, ALB, Z500 hPa, and 3D temperature, water vapor, relative humidity, and wind speeds, were obtained from MERRA-2 via [https://disc.gsfc.nasa.gov/datasets/M2TMNXGLC\\_V5.12.4/summary?keywords=%22MERRA-2%22](https://disc.gsfc.nasa.gov/datasets/M2TMNXGLC_V5.12.4/summary?keywords=%22MERRA-2%22). All data needed to evaluate the conclusions in the paper are presented in Mendeley data (Takano, 2020, doi:<https://doi.org/10.17632/7kxwv8mtbp.2>).

## Acknowledgments

This study was supported by California Department of Water Resources (Award No. 4600010378 AM-30), the U.S. National Oceanic and Atmospheric Administration (NOAA) grant NA19OAR4310243, and the NSF grant AGS-2103820. Yu Gu acknowledges the support by (while serving at) the National Science Foundation. The authors thank the GLDAS, KNMI, CPC, MERRA-2, and CMC for providing datasets that made this work possible. We acknowledge the support of the Joint Institute for Regional Earth System Science and Engineering in the University of California Los Angeles. The computation in this work was completed on the Cheyenne (<https://doi.org/10.5065/D6RX99HX>) cluster system of NCAR's Computational and Information Systems Laboratory, which is sponsored by the National Science Foundation. All data needed to evaluate the conclusions in the paper are presented in Mendeley data (<https://doi.org/10.17632/7kxwv8mtbp.2>).

## References

- Ban, Z., Das, T., Cayan, D., Xiao, M., & Lettenmaier, D. P. (2020). Understanding the asymmetry of annual streamflow responses to seasonal warming in the western United States. *Water Resources Research*, *56*, e2019JA027531. <https://doi.org/10.1029/2020WR027158>
- Burgdorf, A. M., Brönnimann, S., & Franke, J. r. (2019). Two types of North American droughts related to different atmospheric circulation patterns. *Climate of the Past*, *15*, 2053–2065. <https://doi.org/10.5194/cp-15-2053-2019>
- Dai, Y., Zeng, X., Dickinson, R. E., Baker, I., Bonan, G. B., Bosilovich, M. G., et al. (2003). The common land model. *Bulletin of the American Meteorological Society*, *84*(8), 1013–1024. <https://doi.org/10.1175/bams-84-8-1013>
- Das, T., Pierce, D. W., Cayan, D. R., Vano, J. A., & Lettenmaier, D. P. (2011). The importance of warm season warming to western U.S. streamflow changes. *Geophysical Research Letters*, *38*, L23403. <https://doi.org/10.1029/2011GL049660>
- Dawadi, S., & Ahmad, S. (2013). Evaluating the impact of demand-side management on water resources under changing climatic conditions and increasing population. *Journal of Environmental Management*, *114*, 261–275. <https://doi.org/10.1016/j.jenvman.2012.10.015>
- Engström, A., & Ekman, A. M. L. (2010). Impact of meteorological factors on the correlation between aerosol optical depth and cloud fraction. *Geophysical Research Letters*, *37*, L18814. <https://doi.org/10.1029/2010GL044361>
- Francis, J. A., & Vavrus, S. J. (2015). Evidence for a Wavier jet stream in response to rapid Arctic warming. *Environmental Research Letters*, *10*, 014005. <https://doi.org/10.1088/1748-9326/10/1/014005>
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., et al. (2017). The Modern-Era retrospective analysis for research and applications, Version 2 (MERRA-2). *Journal of Climate*, *30*(14), 5419–5454. <https://doi.org/10.1175/JCLI-D-16-0758.1>
- Gleick, P. H. (1986). Methods for evaluating the regional hydrologic impacts of global climatic changes. *Journal of Hydrology*, *88*, 97–116. [https://doi.org/10.1016/0022-1694\(86\)90199-X](https://doi.org/10.1016/0022-1694(86)90199-X)
- Halimi, M., Rezaei, M., Mohammadi, C., & Farajzadeh, M. (2017). Association between cloudiness and rainfall over Fars province in Iran. *Russian Meteorology and Hydrology*, *42*, 671–676. <https://doi.org/10.3103/S1068373917100077>
- Hamlet, A. F., Mote, P. W., Clark, M. P., & Lettenmaier, D. P. (2007). Twentieth-century trends in runoff, evapotranspiration, and soil moisture in the western United States. *Journal of Climate*, *20*, 1468–1486. <https://doi.org/10.1175/JCLI4051.1>
- Hardle, W. K., & Simar, L. (2015). *Applied multivariate statistical analysis* (4th ed.). Berlin, Germany: Springer.
- Hobbins, M., & Barsugli, J. (2020). Threatening the vigor of the Colorado river. *Science*, *367*, 1192–1193. <https://doi.org/10.1126/science.abb3624>
- Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., et al. (2019). High mountain areas, IPCC special report on the ocean and cryosphere in a changing climate (Rep.). Genf, Switzerland: IPCC.
- Hoerling, M., Barsugli, J., Livneh, B., Eischeid, J., Quan, X., & Badger, A. (2019). Causes for the century-long decline in Colorado river flow. *Journal of Climate*, *32*, 8181–8203. <https://doi.org/10.1175/JCLI-D-19-0207.1>
- Huntington, T. G., & Billmire, M. (2014). Trends in precipitation, runoff, and evapotranspiration for rivers draining to the Gulf of Maine in the United States. *Journal of Hydrometeorology*, *15*, 726–743. <https://doi.org/10.1175/JHM-D-13-018.1>
- Johnson, R. A., & Wichern, D. W. (2007). *Applied multivariate statistical analysis* (6th ed.). Hoboken, NJ: Pearson Education Inc.
- Karl, T. R., & Riebsame, W. E. (1989). The impact of decadal fluctuations in mean precipitation and temperature on runoff: A sensitivity study over the United States. *Climate Change*, *15*, 423–447. <https://doi.org/10.1007/BF00240466>
- Kirk, J. P., & Schmidlin, T. W. (2018). Moisture transport associated with large precipitation events in the Upper Colorado River Basin. *International Journal of Climatology*, *38*, 5323–5338. <https://doi.org/10.1002/joc.5734>
- Kirk, J. P., Sheridan, S. C., & Schmidlin, T. W. (2017). Synoptic climatology of the early 21st century drought in the Colorado River Basin and relationships to reservoir water levels. *International Journal of Climatology*, *37*, 2424–2437. <https://doi.org/10.1002/joc.4855>
- Lamquin, N., Stubenrauch, C. J., & Pelon, J. (2009). Upper tropospheric humidity and cirrus geometrical and optical thickness: Relationships inferred from 1 year of collocated AIRS and CALIPSO data. *Journal of Geophysical Research*, *114*, D00A08. <https://doi.org/10.1029/2008JD010012>

- Lehner, F., Deser, C., Simpson, I. R., & Terray, L. (2018). Attributing the U.S. Southwest's recent shift into drier conditions. *Geophysical Research Letters*, *45*, 6251–6261. <https://doi.org/10.1029/2018GL078312>
- Liu, C., Ikeda, K., Rasmussen, R., Barlage, M., Newman, A. J., Prein, A. F., et al. (2017). Continental-scale convection-permitting modeling of the current and future climate of North America. *Climate Dynamics*, *49*, 71–95. <https://doi.org/10.1007/s00382-016-3327-9>
- Mavromatis, T., & Stathis, D. (2011). Response of the water balance in Greece to temperature and precipitation trends. *Theoretical and Applied Climatology*, *104*, 13–24. <https://doi.org/10.1007/s00704-010-0320-9>
- McCabe, G. J., Wolock, D. M., Pederson, G. T., Woodhouse, C. A., & McAfee, S. (2017). Evidence that recent warming is reducing upper Colorado river flows. *Earth Interactions*, *21*, 1–14. <https://doi.org/10.1175/EI-D-17-0007.1>
- Miller, W. P., & Piechota, T. C. (2008). Regional analysis of trend and step changes observed in hydroclimatic variables around the Colorado river basin. *Journal of Hydrometeorology*, *9*, 1020–1034. <https://doi.org/10.1175/2008JHM988.1>
- Milly, P. C. D., & Dunne, K. A. (2020). Colorado River flow dwindles as warming-driven loss of reflective snow energizes evaporation. *Science*, *367*, 1252–1255. <https://doi.org/10.1126/science.aay9187>
- Milly, P. C. D., Kam, J., & Dunne, K. A. (2018). Sensitivity of annual streamflow to air temperature. *Water Resources Research*, *54*, 2624–2641. <https://doi.org/10.1002/2017WR021970>
- Mishra, A. K. (2019). Investigating changes in cloud cover using the long-term record of precipitation extremes. *Meteorological Applications*, *26*, 108–116. <https://doi.org/10.1002/met.1745>
- Namias, J. (1983). Some causes of United States drought. *Journal of Applied Meteorology and Climatology*, *22*, 30–39. [https://doi.org/10.1175/1520-0450\(1983\)022<0030:SCOUSD>2.0.CO;2](https://doi.org/10.1175/1520-0450(1983)022<0030:SCOUSD>2.0.CO;2)
- Nash, L. L., & Gleick, P. H. (1991). Sensitivity of streamflow in the Colorado Basin to climatic changes. *Journal of Hydrology*, *125*, 221–241. [https://doi.org/10.1016/0022-1694\(91\)90030-L](https://doi.org/10.1016/0022-1694(91)90030-L)
- Nowak, K., Hoerling, M., Rajagopalan, B., & Zagona, E. (2012). Colorado river basin hydroclimatic variability. *Journal of Climate*, *25*, 4389–4403. <https://doi.org/10.1175/JCLI-D-11-00406.1>
- Painter, T. H., Deems, J. S., Belnap, J., Hamlet, A. F., Landry, C. C., & Udall, B. (2010). Response of Colorado River runoff to dust radiative forcing in snow. *Proceedings of the National Academy of Sciences of the United States of America*, *107*, 17125–17130. <https://doi.org/10.1073/pnas.0913139107>
- Pascolini-Campbell, M., Reager, J. T., Chandanpurkar, H. A., & Rodell, M. (2021). A 10 per cent increase in global land evapotranspiration from 2003 to 2019. *Nature*, *593*, 543–547. <https://doi.org/10.1038/s41586-021-03503-5>
- Sakakibara, H. (1978). *Statistical relations between precipitation formation and quantities of synoptic field*. [https://doi.org/10.2151/jmsj1965.56.3\\_211](https://doi.org/10.2151/jmsj1965.56.3_211)
- Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., et al. (2010). Investigating soil moisture-climate interactions in a changing climate: A review. *Earth-Science Reviews*, *99*, 125–161. <https://doi.org/10.1016/j.earscirev.2010.02.004>
- Smith, R. A., & Kummerow, C. D. (2013). A comparison of in situ, reanalysis, and satellite water budgets over the Upper Colorado River Basin. *Journal of Hydrometeorology*, *14*, 888–905. <https://doi.org/10.1175/JHM-D-12-0119.1>
- Takano, Y. (2020). *Cloud influence on UCRB runoff, Mendeley data*, V2. <https://doi.org/10.17632/7kxwv8mtbp.2>
- Udall, B., & Overpeck, J. (2017). The twenty-first century Colorado River hot drought and implications for the future. *Water Resources Research*, *53*, 2404–2418. <https://doi.org/10.1002/2016WR019638>
- Vano, J. A., Das, T., & Lettenmaier, D. P. (2012). Hydrologic sensitivities of Colorado River runoff to changes in precipitation and temperature. *Journal of Hydrometeorology*, *13*, 932–949. <https://doi.org/10.1175/JHM-D-11-069.1>
- Vano, J. A., & Lettenmaier, D. P. (2014). A sensitivity-based approach to evaluating future changes in Colorado River discharge. *Climate Change*, *122*, 621–634. <https://doi.org/10.1007/s10584-013-1023-x>
- Vano, J. A., Nijssen, B., & Lettenmaier, D. P. (2015). Seasonal hydrologic responses to climate change in the Pacific Northwest. *Water Resources Research*, *51*, 1959–1976. <https://doi.org/10.1002/2014WR015909>
- Walcek, C. J. (1994). Cloud cover and its relationship to relative humidity during a springtime midlatitude cyclone. *Monthly Weather Review*, *122*, 1021–1035. [https://doi.org/10.1175/1520-0493\(1994\)122<1021:CCAIRT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<1021:CCAIRT>2.0.CO;2)
- Wigley, T. M. L., & Jones, P. D. (1985). Influences of precipitation changes and direct CO<sub>2</sub> effects on streamflow. *Nature*, *314*(6007), 149–152. <https://doi.org/10.1038/314149a0>
- Woodhouse, C. A., Pederson, G. T., Morino, K., McAfee, S. A., & McCabe, G. J. (2016). Increasing influence of air temperature on Upper Colorado River streamflow. *Geophysical Research Letters*, *43*, 2174–2181. <https://doi.org/10.1002/2015GL067613>
- Xiao, M., Udall, B., & Lettenmaier, D. P. (2018). On the causes of declining Colorado River streamflows. *Water Resources Research*, *54*, 6739–6756. <https://doi.org/10.1029/2018WR023153>
- Yue, S., & C. Wang (2004). The Mann-Kendall test modified by effective sample size to detect trend in serially correlated hydrological series. Paper presented at Water Resources Management.
- Zaitchik, B. F., Rodell, M., & Olivera, F. (2010). Evaluation of the Global Land Data Assimilation System using global river discharge data and a source-to-sink routing scheme. *Water Resources Research*, *46*, W06507. <https://doi.org/10.1029/2009WR007811>
- Zhao, B., Gu, Y., Liou, K. N., Wang, Y., Liu, X., Huang, L., et al. (2018). Type-dependent responses of ice cloud properties to aerosols from satellite retrievals. *Geophysical Research Letters*, *45*, 3297–3306. <https://doi.org/10.1002/2018GL077261>
- Zhao, S., Fu, R., Zhuang, Y., & Wang, G. (2021). Long-lead seasonal prediction of streamflow over the upper Colorado river basin: The role of the Pacific sea surface temperature and beyond. *Journal of Climate*, *34*, 6855–6873. <https://doi.org/10.1175/JCLI-D-20-0824.1>