



Reply to comment by Roberto Rondanelli and Richard S. Lindzen on “Variations in convective precipitation fraction and stratiform area with sea surface temperature”

Hui Su,¹ Jonathan H. Jiang,¹ J. David Neelin,² Brian H. Kahn,¹ Joe W. Waters,¹ Nathaniel J. Livesey,¹ and Yu Gu²

Received 22 July 2009; revised 21 December 2009; accepted 30 December 2009; published 18 March 2010.

Citation: Su, H., J. H. Jiang, J. D. Neelin, B. H. Kahn, J. W. Waters, N. J. Livesey, and Y. Gu (2010), Reply to comment by Roberto Rondanelli and Richard S. Lindzen on “Variations in convective precipitation fraction and stratiform area with sea surface temperature,” *J. Geophys. Res.*, 115, D06203, doi:10.1029/2009JD012872.

[1] *Lindzen et al.* [2001] reported that cirrus anvil cloud fraction normalized by deep convective core cloud fraction decreases with cloud-weighted sea surface temperature (SST) by about $-20\%/K$, while *Su et al.* [2008] showed tropical-mean upper tropospheric (UT) cloud fraction normalized by precipitation decreases with cloud-weighted SST more slowly at around $-2\%/K$. Throughout the text, we use “ $\%/K$ ” to indicate the relative change of normalized cloud fraction per degree change of SST. In their comments to *Su et al.* [2008], *Rondanelli and Lindzen* [2009] (hereafter RL09) examined several methodological choices that may contribute to the reported fractional decreases. First, they presented a regression analysis of normalized anvil cloud fraction onto SST using binned cloud fraction data within $0.5^{\circ}C$ of SST and $1^{\circ} \times 1^{\circ}$ gridded data. Both binned and gridded regressions show that anvil cloud fraction normalized by convective core cloud fraction decreases with SST at a rate of approximately $-20\%/K$. In their analysis, cirrus anvil fraction is based on brightness temperature (BT) from the Tropical Rainfall Measuring Mission (TRMM) Visible and Infrared Scanner (VIRS) with values between 220 K and 260 K. Their convective core cloud fraction, which served as a measure of convective mass flux, is based on lower brightness temperature ($BT < 220$ K). The normalization focuses on the relative change of anvil cloud fraction with respect to unit convective mass flux, i.e., the efficiency of anvil formation due to convective detrainment. *Su et al.* [2008] used precipitation to normalize high-altitude cloud fraction to account for the competition between precipitation and anvil formation for a given moisture supply. When RL09 applied to their data the same methodology as in the study by *Su et al.* [2008] which regressed the daily tropical-averaged cloud fraction onto the cloud-weighted SST, a decrease of $-6\%/K$ was obtained, which was rather close in magnitude to the finding of *Su et al.* [2008]. This shows that different methodologies contribute to the differences in the rates of relative changes.

[2] Here, we further test if different data sets could cause the differences in the rates of relative change. We replicate the analysis by RL09’s Figure 1 with the $1^{\circ} \times 1^{\circ}$ gridded cloud fraction data from the Aqua Atmospheric Infrared Sounder (AIRS) normalized by the TRMM precipitation. The daily AIRS UT (pressure < 300 hPa) cloud fraction normalized by precipitation shows a relative change of $-18\%/K$ based on the decaying exponential fit to all daily data from January to March (the same months as used by RL09) 2006, while the binned cloud fraction shows a relative change of $-25\%/K$ (Figure 1). Both rates are similar to the counterparts from RL09. The correlation between the precipitation-normalized cloud fraction and SST is only -0.05 . The rates of change are rather sensitive to analysis periods and data uncertainties, but the sign is always negative. Therefore, a decrease of *normalized* cirrus fraction with increasing SST appears to be robust, although different analysis methodologies can give different magnitude of relative change.

[3] However, we note that the UT cloud fraction is not simply proportional to precipitation. This has significant implications for the rate of change of precipitation-normalized cloud fraction with SST. Figure 2 illustrates the relationships among tropical UT cloud fraction, precipitation and SST using the $1^{\circ} \times 1^{\circ}$ gridded data from Figure 1. Similar relations for tropical-mean values have been shown by *Su et al.* [2008]. Precipitation, binned in $0.5^{\circ}C$ SST intervals, increases approximately linearly with SST (Figure 2a). Cloud fraction increases with SST $> 27^{\circ}C$, but decreases with SST $< 27^{\circ}C$ (Figure 2b), as the two regimes are associated with different large-scale circulation patterns. Cloud fraction increases with precipitation (Figure 2c), rather rapidly for small values of precipitation and less so when precipitation $> \sim 20$ mm/d. This “saturation” behavior of cloud fraction with precipitation may contribute partly to the negative slope of precipitation-normalized cloud fraction versus SST. On the other hand, we notice that there is a high occurrence of large cloud fraction when precipitation is zero or very small. Thus, if one performs a linear least squares fit to the full range of cloud fraction versus precipitation, one obtains a nonzero intercept, corresponding to the fact that some high clouds are present regardless of precipitation. These high clouds may not be of convective origin and complicate the relation between the precipitation-normalized cloud fraction and SST.

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA.

²Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA.





Figure 1. Scatterplots of the upper tropospheric cloud fraction (with cloud top pressure less than 300 hPa) normalized by precipitation versus sea surface temperature (similar to Figure 1 of *Rondanelli and Lindzen* [2009] but with the data used by *Su et al.* [2008]). The data are from January to March 2006 for oceanic regions within 15°S–15°N. The gray crosses represent individual 1° × 1° gridded daily AIRS UT cloud fraction normalized by TRMM precipitation (greater than 0.02 mm/d). The black dots are the binned normalized cloud fraction for 0.5°C SST bins. The solid black curve is the decaying exponential fit to the binned data, and the dashed black curve is the decaying exponential fit to all individual gridded data. “Rate” refers to the relative change of precipitation-normalized cloud fraction versus SST based on the decaying exponential fit.

[4] Using the linear fit between cloud fraction (CFR) and precipitation (P) (which rather poorly captures their relationship) to illustrate the importance of the nonzero intercept, we can write $CFR(SST) = \alpha(SST)P(SST) + \beta(SST)$, where $\alpha(SST)$ represents the anvil formation efficiency and $\beta(SST)$ denotes the nonzero intercept, both of which may vary with SST. Hence, the rate of change for precipitation-normalized cloud fraction with SST can be expressed as $\frac{d(CFR/P)}{dSST} = \frac{d\alpha}{dSST} + \frac{1}{P} \frac{d\beta}{dSST} - \frac{\beta}{P^2} \frac{dP}{dSST}$. The term $\frac{d\alpha}{dSST}$ is of interest for cloud-climate feedback. However, only when β is zero can we reliably obtain $\frac{d\alpha}{dSST}$ directly from $\frac{d(CFR/P)}{dSST}$. When β is not zero and $\frac{dP}{dSST}$ is positive, the $-\frac{\beta}{P^2} \frac{dP}{dSST}$ term will make a significant contribution to the rate of relative change of precipitation-normalized cloud fraction versus SST. In the

Figure 2. (a) The 1° × 1° gridded precipitation (15°S–15°N) binned on 0.5°C SST bins. The dashed line is the least squares linear fit to the binned data, with the regression equation shown. (b) The 1° × 1° gridded UT cloud fraction binned in 0.5°C SST intervals. (c) The gridded UT cloud fraction as a function of gridded precipitation. The color shading is the joint density with high density in red and low density in blue. The black dots are cloud fraction binned in 2 mm/d precipitation intervals and the solid line is the least squares linear fit to the full range of data.

data we used, the term $-\frac{\beta}{P^2} \frac{dP}{dSST}$ would make a rate of relative change about $-14\%/K$ for $P = 10$ mm/d and $-6\%/K$ for $P = 20$ mm/d, a significant contribution to the rates depicted in Figure 1. Thus, it is not clear whether the anvil formation efficiency α increases or decreases with SST, given unknown $\frac{d\beta}{dSST}$ and other uncertainties associated



with the data. Hence, while normalization by some measure of the convective source is desirable, the nonproportionality between cloud fraction and precipitation appears to compromise the usefulness of a normalization procedure that assumes proportionality (such as dividing the anvil cloud fraction by precipitation). We expect that similar nonproportionality applies to the anvil cloud fraction and convective core fractions as defined by BT used by RL09 (R. Rondanelli, personal communication, 2009). Thus similar uncertainty exists when interpreting the rates of relative change from their analysis.

[5] Cloud fraction, of course, is only one cloud property that is relevant to the radiative effects of clouds. Other cloud properties, such as cloud water content (as shown by *Su et al.* [2008]), cloud optical depth, cloud particle size and habits, are all important to the net cloud radiative forcing. The magnitude and sign of cloud climate feedback requires further study, with sustained (decadal or longer) simultaneous observations of multiple cloud parameters.

[6] **Acknowledgment.** This work was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

References

- Lindzen, R. S., M.-D. Chou, and A. Y. Hou (2001), Does the Earth have an adaptive infrared iris?, *Bull. Am. Meteorol. Soc.*, *82*, 417–432, doi:10.1175/1520-0477(2001)082<0417:DTEHAA>2.3.CO;2.
- Rondanelli, R., and R. S. Lindzen (2009), Comment on “Variations of tropical upper tropospheric clouds with sea surface temperature and implications for radiative effects” by H. Su et al., *J. Geophys. Res.*, *115*, D06202, doi:10.1029/2008JD011189.
- Su, H., et al. (2008), Variations of tropical upper tropospheric clouds with sea surface temperature and implications for radiative effects, *J. Geophys. Res.*, *113*, D10211, doi:10.1029/2007JD009624.
-
- Y. Gu and J. D. Neelin, Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA 90095, USA.
- J. H. Jiang, B. H. Kahn, N. J. Livesey, H. Su, and J. W. Waters, Jet Propulsion Laboratory, California Institute of Technology, M/S 183-701, 4800 Oak Grove Dr., Pasadena, CA 91109-8099, USA. (Hui.Su@jpl.nasa.gov)